

Figures on BNT162b2 effectiveness and safety at age 12-15 years


Figures:

1. Translated Israel figures on myocarditis following BNT162b2
2. Miron et al analysis on BNT162b2 effectiveness



Preliminary summary:

The 1st dose of BNT162b2 was not significantly associated with myocarditis and was effective against COVID-19 before the increase in the Delta variant, which may reduce 1st dose effectiveness. The 2nd dose of BNT162b2 was associated with myocarditis, but cases are rare and mostly mild, and since 2nd dose likely improves the effectiveness against Delta, it could justify the myocarditis risk. If the UK administers dose-1 to ages 12-15 years, there will be another 21 days until 2nd dose administration to examine the dose 1 and dose 2 effectiveness against the Delta variant, using additional data from the UK, US and Israel.


1. Translated figures on myocarditis & BNT162b2



1. Translated figures on myocarditis & BNT162b2



1. Translated figures on myocarditis & BNT162b2



1. Translated figures on myocarditis & BNT162b2

Calculation of myocarditis risk in all vaccinated
חישוב הסיכון למיוקרדיטיס בכלל המתחסנים

Dose 2 Case per ... vaccinated מנה שניה מקרה אחד ל... מתחסנים	Dose 1 Case per ... vaccinated מנה ראשונה מקרה אחד ל... מתחסנים	Age and sex groups קבוצת גיל ומין	
6,230	90,511	Male זכר	16-19
No reported cases in this group לא דווחו מקרים בקבוצה זו	No reported cases in this group לא דווחו מקרים בקבוצה זו	Female נקבה	
10,463	132,724	Male זכר	20-24
58,385	No reported cases in this group לא דווחו מקרים בקבוצה זו	Female נקבה	
25,304	No reported cases in this group לא דווחו מקרים בקבוצה זו	Male זכר	25-29
218,014	No reported cases in this group לא דווחו מקרים בקבוצה זו	Female נקבה	
71,785	393,9418	Male זכר	30+

1. Translated figures on myocarditis & BNT162b2

נכון ל-03/08/2021 דווחו 5 מקרים של דלקת בשריר הלב בגילאי 12-15 מתוך 201,540 מקבלי מנה ראשונה ו-153,491 מקבלי מנה שניה.

As of 03/08/2021, 5 cases were reported of myocarditis at ages 12-15 years, out of 201,540 1st doses and 153,491 2nd doses

מנה שניה:

2nd dose

4 מקרים, כולם זכרים

4 case, male

3-5 ימים לאחר קבלת החיסון

3-5 days after vaccine

כולם שוחררו להמשך טיפול בקהילה

Released for community care

מנה ראשונה:

1st dose

• מקרה 1, זכר.

1 case male


• 5 ימים לאחר קבלת החיסון

5 days after vaccine


• שוחררו להמשך טיפול בקהילה

Released for community care

2. Reanalysis of BNT162b2 age 12-15 years



2. Reanalysis of BNT162b2 RCT age 12-15 years



2. Reanalysis of BNT162b2 RCT age >15 years

